congenital bilateral absence of the vas deferens

Congenital bilateral absence of the vas deferens occurs in males when the tubes that carry sperm out of the testes (the vas deferens) fail to develop properly. Although the testes usually develop and function normally, sperm cannot be transported through the vas deferens to become part of semen. As a result, men with this condition are unable to father children (infertile) unless they use assisted reproductive technologies. This condition has not been reported to affect sex drive or sexual performance.

This condition can occur alone or as a sign of cystic fibrosis, an inherited disease of the mucus glands. Cystic fibrosis causes progressive damage to the respiratory system and chronic digestive system problems. Many men with congenital bilateral absence of the vas deferens do not have the other characteristic features of cystic fibrosis; however, some men with this condition may experience mild respiratory or digestive problems.

Frequency

This condition is responsible for 1 percent to 2 percent of all infertility in men.

Genetic Changes

Mutations in the CFTR gene cause congenital bilateral absence of the vas deferens.

More than half of all men with this condition have mutations in the *CFTR* gene. Mutations in this gene also cause cystic fibrosis. When congenital bilateral absence of the vas deferens occurs with *CFTR* mutations, it is considered a form of atypical cystic fibrosis.

The protein made from the *CFTR* gene forms a channel that transports negatively charged particles called chloride ions into and out of cells. The flow of chloride ions helps control the movement of water in tissues, which is necessary for the production of thin, freely flowing mucus. Mucus is a slippery substance that lubricates and protects the linings of the airways, digestive system, reproductive system, and other organs and tissues.

Mutations in the *CFTR* gene disrupt the function of the chloride channels, preventing them from regulating the flow of chloride ions and water across cell membranes. As a result, cells in the male genital tract produce mucus that is abnormally thick and sticky. This mucus clogs the vas deferens as they are forming, causing them to deteriorate before birth.

In instances of congenital bilateral absence of the vas deferens without a mutation in the *CFTR* gene, the cause of this condition is often unknown. Some cases are associated with other structural problems of the urinary tract.

Inheritance Pattern

When this condition is caused by mutations in the *CFTR* gene, it is inherited in an autosomal recessive pattern. This pattern of inheritance means that both copies of the gene in each cell have a mutation. Men with this condition who choose to father children through assisted reproduction have an increased risk of having a child with cystic fibrosis. If congenital absence of the vas deferens is not caused by mutations in *CFTR*, the risk of having children with cystic fibrosis is not increased.

Other Names for This Condition

- Absence of vas deferens
- Absent vasa
- CAVD
- CBAVD
- Congenital absence of vas deferens
- Congenital aplasia of vas deferens
- congenital bilateral absence of vas deferens

Diagnosis & Management

These resources address the diagnosis or management of congenital bilateral absence of the vas deferens:

- GeneReview: Cystic Fibrosis and Congenital Absence of the Vas Deferens https://www.ncbi.nlm.nih.gov/books/NBK1250
- Genetic Testing Registry: Congenital bilateral absence of the vas deferens https://www.ncbi.nlm.nih.gov/gtr/conditions/C0403814/
- MedlinePlus Encyclopedia: Infertility https://medlineplus.gov/ency/article/001191.htm
- MedlinePlus Encyclopedia: Pathway of sperm (image) https://medlineplus.gov/ency/imagepages/19073.htm
- MedlinePlus Health Topic: Assisted Reproductive Technology https://medlineplus.gov/assistedreproductivetechnology.html

These resources from MedlinePlus offer information about the diagnosis and management of various health conditions:

- Diagnostic Tests
 https://medlineplus.gov/diagnostictests.html
- Drug Therapy https://medlineplus.gov/drugtherapy.html

- Surgery and Rehabilitation https://medlineplus.gov/surgeryandrehabilitation.html
- Genetic Counseling https://medlineplus.gov/geneticcounseling.html
- Palliative Care https://medlineplus.gov/palliativecare.html

Additional Information & Resources

MedlinePlus

- Encyclopedia: Infertility https://medlineplus.gov/ency/article/001191.htm
- Encyclopedia: Pathway of sperm (image)
 https://medlineplus.gov/ency/imagepages/19073.htm
- Health Topic: Assisted Reproductive Technology https://medlineplus.gov/assistedreproductivetechnology.html
- Health Topic: Male Infertility https://medlineplus.gov/maleinfertility.html

Genetic and Rare Diseases Information Center

 Congenital bilateral absence of the vas deferens https://rarediseases.info.nih.gov/diseases/5461/congenital-bilateral-absence-of-the-vas-deferens

Educational Resources

- American Society for Reproductive Medicine: Male Infertility http://www.asrm.org/topics/detail.aspx?id=1331
- MalaCards: congenital bilateral absence of vas deferens http://www.malacards.org/card/congenital_bilateral_absence_of_vas_deferens
- Orphanet: Congenital bilateral absence of vas deferens http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=48

Patient Support and Advocacy Resources

- Canadian Cystic Fibrosis Foundation http://www.cysticfibrosis.ca/
- Path2Parenthood http://www.path2parenthood.org/article/its-in-the-male/
- Resolve: The National Infertility Association http://www.resolve.org

GeneReviews

 Cystic Fibrosis and Congenital Absence of the Vas Deferens https://www.ncbi.nlm.nih.gov/books/NBK1250

Genetic Testing Registry

 Congenital bilateral absence of the vas deferens https://www.ncbi.nlm.nih.gov/gtr/conditions/C0403814/

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28absence+of+vas+deferens %5BTIAB%5D%29+OR+%28absent+vasa%5BTIAB%5D%29+OR+%28cavd%5BTIAB%5D%29+OR+%28congenital+absence+of+vas+deferens%5BTIAB%5D%29+OR+%28congenital+aplasia+of+vas+deferens %5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D

OMIM

 VAS DEFERENS, CONGENITAL BILATERAL APLASIA OF http://omim.org/entry/277180

Sources for This Summary

- Cuppens H, Cassiman JJ. CFTR mutations and polymorphisms in male infertility. Int J Androl. 2004 Oct;27(5):251-6. Review.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15379964
- Daudin M, Bieth E, Bujan L, Massat G, Pontonnier F, Mieusset R. Congenital bilateral absence
 of the vas deferens: clinical characteristics, biological parameters, cystic fibrosis transmembrane
 conductance regulator gene mutations, and implications for genetic counseling. Fertil Steril. 2000
 Dec;74(6):1164-74.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/11119745
- GeneReview: Cystic Fibrosis and Congenital Absence of the Vas Deferens https://www.ncbi.nlm.nih.gov/books/NBK1250
- Gilljam M, Moltyaner Y, Downey GP, Devlin R, Durie P, Cantin AM, Zielenski J, Tullis DE. Airway inflammation and infection in congenital bilateral absence of the vas deferens. Am J Respir Crit Care Med. 2004 Jan 15;169(2):174-9. Epub 2003 Oct 9.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/14551163
- Jarzabek K, Zbucka M, Pepinski W, Szamatowicz J, Domitrz J, Janica J, Wolczynski S, Szamatowicz M. Cystic fibrosis as a cause of infertility. Reprod Biol. 2004 Jul;4(2):119-29. Review. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15297887
- Nick JA, Rodman DM. Manifestations of cystic fibrosis diagnosed in adulthood. Curr Opin Pulm Med. 2005 Nov;11(6):513-8. Review.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/16217177

 Noone PG, Knowles MR. 'CFTR-opathies': disease phenotypes associated with cystic fibrosis transmembrane regulator gene mutations. Respir Res. 2001;2(6):328-32. Epub 2001 Aug 9. Review.

Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/11737931
Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC64805/

 Samli H, Samli MM, Yilmaz E, Imirzalioglu N. Clinical, andrological and genetic characteristics of patients with congenital bilateral absence of vas deferens (CBAVD). Arch Androl. 2006 Nov-Dec; 52(6):471-7.

Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/17050329

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/congenital-bilateral-absence-of-the-vas-deferens

Reviewed: January 2008 Published: February 7, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services